



# Late Holocene sedimentation in Iberian Range karstic lakes: Facies model, depositional evolution and climatic implications

F. Barreiro-Lostres \* Pyrenean Institute of Ecology, CSIC, Spain  
A. Moreno Pyrenean Institute of Ecology, CSIC, Spain  
S. Giralt Earth Sciences Jaume Almera Institute, CSIC, Spain  
P. Mata Spanish Geological Survey (IGME), Spain  
J. Aranbarri Pyrenean Institute of Ecology, CSIC, Spain  
B. Valero-Garcés Pyrenean Institute of Ecology, CSIC, Spain






# Mediterranean Records in karstic terrains...

also small lakes and short sequences

- The potential of small karstic lakes
- Facies as paleohydrological tools
- Climate and Human interactions during the last millennia



# The Iberian Peninsula:climates





# Lacustrine Records



B. DE LA MORA

ARREO

TARAVILLA

ZOÑAR

MONTCORTÉS

ESTANYA

Minorca contourite

Tagus prodelta


Montcortès Corella et al.  
(S12-P02)

Banyoles Morellón et al.  
(S12-P05)

Basa Mora Moreno et al.  
(S12-P09)

# Zoñar Lake (southern Spain)

Martín-Puertas et al., (2008, 2009, 2010)




- Arid period prior to 2.8 BP
- Iberian – Roman Period → humid (2.5 – 1.5 ka BP)  
(with a dry phase around 2 ka BP)
- Arid MCA
- Humid LIA





# The Pyrenees: Estanya Lake

Morellón et al.,(2010)




- Some dry phases during Roman times
- Arid MCA
- Humid LIA but with several phases



# The Pyrenees: Montcortès Lake

Corella et al. (S12-P02)




- A varved record for the last 1.5 ka!
- Arid MCA
- Cold (and humid) LIA




# The Iberian Range: La Cruz

Julià et al., (1998), Romero-Viana, 2010)



- 1.75 m long core:
  - A varved record since the LIA
  - Arid MWA
  - Cold (and humid) LIA




Reconstructed precipitation  
for the last 400 yr based on  
calcite laminae thickness



## Introduction & Location

# Last 2 ka


- The IRHP (2600-1600 yr BP) is the most humid period in the last 3500 yr (Zoñar lake) but not so much in northern Spain
- The MCA (IX- XV centuries): a relatively arid period in all records
- The LIA (1300-1850 yr AD) relatively wetter
- **Regional (latitudinal) variability?**
- **Higher resolution (varves?)**



(Morellón et al., in press)



# In search of varved sediments...

- Do Iberian karstic lake sequences contain high resolution paleoclimate information for the last millennia?..





# The Iberian Peninsula : geology



# Large carbonate areas affected by endo- and exokarstic processes...

... developing


-travertine-damned lakes

-circular funnel-shaped depressions

-These systems are sensitive to climate changes and their **sedimentary records** are a good approach to study the fluctuations of the **hydrological cycle**








- Lake hydrology:

→ Groundwater input vs. evaporation output


- Evidences of relatively rapid response of water level to rainfall variability:

- Dry year 1995 → isolation of NW and SE sub-basins

- Humid year 2003 → sill separating the 2 sub-basins submerged, at 2-3 m water depth



# Lakes in the Iberian Range







## La Parra Lake



Located on a doline system developed on Cretaceous carbonatic formations.

# The Lake




- **Inputs:** Groundwater and runoff-water.
- **Outputs:** Irrigation wells

The lake is **Holomictic**







# The Cores



# Sedimentology


Detrital Facies		Description
A	<i>Grey fine silts</i>	Massive, in beds of 10-25 cm and diffuse contacts









# Sedimentology

Detrital Facies		Description
B	<i>Black fine silts</i>	Massive in isolated levels of 2-5 cm with net boundaries
Detrital Facies		Description
C	<i>Medium brown silts</i>	Layers of 5-20 cm with difusse boundaries.



# Sedimentology

Detrital Facies		Description
D,E	<i>Coarse dark &amp; clear grey silts</i>	Massive levels of 5-10 cm with irregular basal boundaries and abundant OM fragments

**Facies D****Facies E**



# Sedimentology

	Detrital Facies	Description
F	<i>Very fine green sands</i>	Massive, forming isolated levels of 3-4 cm in a green silty matrix
G	<i>Medium brown sands</i>	Massive, in levels of 2-10 cm with irregular boundaries with a silty fine-medium matrix
H	<i>Coarse brown sands</i>	Massive, with angulous carbonatic pebbles in a level of 10 cm with very irregular boundaries

Facies F



Facies G



Facies H




# Sedimentology

	Detrital Facies	Description
I	<b><i>Fine brown gravels</i></b>	Massive, in a level of 2 cm with irregular boundaries. Pebbles are carbonatic.
J	<b><i>Medium brown gravels</i></b>	Massive, in a level of 30 cm with very irregular boundaries. Pebbles are carbonatic and with a silty matrix.

Facies I



Facies J






10  $\mu$ m

III = 15.0 kV W11 = 0.4 mm Signal = 1.00E+ Signal A = Si 2  
 ESB Grd = 720 V Neg = 1.00 kX Pixel Size = 105.2 nm Signal B = AuB  
 Probe = 170 pA Date: 05/05/2011 File Name = 05\_01b

Scanned with a Zeiss DSM 940 SEM  
 www.zeiss.com

# Unit III: 1<sup>st</sup> lacustrine deposits → Increase of water table



- Laminated facies grading to coarser silts:


- Rapid water table level changes;

- Littoral and shallower environment deposit .





# Unit II: Detrital fine facies



- Increase of fine siliciclastic inputs


- Implies:

- Higher lake levels

- Higher local anthropogenic impacts (erosion).



# Bottom Unit I: Coarser facies deposition




● Coarser facies at the center of the lake:

● Implies shallower levels and an increase on proximal detrital inputs





# Medium Unit I: Fine and Laminated facies




- Higher lake levels.
- Laminated facies (more anoxic conditions at lake bottom)

(similar pattern than nearby La Cruz lake)



# Top Unit I: Dominance of fine massive silts



- Laminated facies disappear


- Relative drop of lake level during recent times






# Age Model

## Age vs Depth cal yr AD




# Geochemical proxies






During the **late IRHP** (300-500 AD), occurs the onset of lacustrine sedimentation, with a rapid increase on detrital input (Ti, Si). Later a drop of water table (400 AD, more clastic facies) occurs.




The **DA** (550-900 AD) start with higher clastic inputs (Ti, Si) to the lake and coarser facies occurs (lower lake levels). Later on, fine facies deposited (higher lake levels).






- During the **onset of the MCA** (900 AD) Ti content is low and coarser carbonate facies with detrital OM are deposited.
- High values of Sr/Ti, Ca/Ti and TOC suggest a shallower lacustrine environment.




During the **LIA** (1300-1850), laminated anoxic facies occur. Sr/Ti, Ca/Ti and TIC have high values around 1500 AD → Higher lake levels (similar pattern La Cruz lake).





# The Lakes







# The Lake




2 sub-basins:





# The Lake




2 sub-basins:

- **Holomictic:** shallower (12 m depth)



# The Lake



2 sub-basins:

- **Holomictic:** shallower (12 m depth)
- **Meromictic:** funnel-shaped (20 m) → Has a permanent saline **anoxic monimolimnion** below 12 m due to groundwater seepage.



# El Tobar



Connection with  
nearby reservoir  
“La Tosca”





# The Lake



Fresh  
water  
springs

Connection with  
nearby reservoir  
“La Tosca”



# The Lake




Fresh  
water  
springs

Diffuse saline  
groundwater

Connection with  
nearby reservoir  
“La Tosca”




# The Lake





# Cores






## Sedimentology

6 main facies have been identified





## Legend

### Facies



E: Greenish grey coarse carbonatic sand



F: Grey clayish silt with pebbles



**Facies E: Greenish grey coarse carbonatic-rich sand with plant and gastropod remains and diatoms.**


**Facies F: Grey clayish silt with pebbles.**






**Facies D: Grey coarse carbonatic silty mud with some Qz, silicates, diatoms and OM. More clastic.**

**Slumps**






## Sedimentology

**Facies C: Black carbonatic mud** with pyrite, diatoms, amorphous OM and plant remains.

There are two calcite groups: i) small crystals (autigenic) and ii) big ones (detrital).






## Sedimentology

**Facies B: Brown fine carbonatic mud with small Qz crystals, some gypsum, diatoms and amorphous OM.**





# El Tobar



## Sedimentology

**Facies A: Finely laminated carbonatic mud** composed by black, brown and grey laminae:

**Black:** carbonatic mud with diatoms.

**Grey:** carbonatic-silty mud with Qz.

**Brown:** carbonatic mud with Qz.




# Chronology... in progress



- Preliminary AMS  $^{14}\text{C}$  dates shows a basal age ca. 100 BC.  $^{137}\text{Cs}$  and some chronological markers suggest high sedimentation rates (1cm/yr)




# Geochemical proxies







During the RP occurs a rapid transition from a very shallow lake (sands with gastropod and silts facies) to a deep and meromictic lake (laminated facies appear)



During Medieval times, there is an increase in coarse silt facies and also an increase in gypsum.




During the LIA , a decrease in clastic facies and a relatively decrease in gypsum occurs.





# The Lakes





# El Tejo




The deepest lake in the region ( 28 m)  
A doline lake similar to La Parra  
The best candidate for varves!







# El Tejo



- About 6 m long lake sequence
- Very thin “varved” intervals
- Bottom coarse, shallower facies and sharp transition to deeper water (banded, faintly laminated facies)
- Preliminary dating: transition about 2000 BP




# To conclude...

## Conclusions



- Iberian Karstic lakes display a large facies variability during the last 2000 years
- Facies and geochemical signatures are valuable paleohydrological (paleoclimate) indicators
- Detail facies models are needed

## Conclusions



In the Iberian Peninsula, the IRHP, the MCA and the LIA had a strong impact on the hydrological cycle




# To conclude...

- Intense dry period prior to the Roman Period ( or within the RP in southern Spain)
- RP:
  - Increase humidity, particularly at the end of the RP
  - A significant latitudinal component exists:  
RP wetter in the S than in the N
- The MCA is a drier period
- The LIA is a wetter period with several internal phases
- Human impact:
  - 2 periods of increased human activity near the lakes (at the end of the RP and during Medieval times)
  - 2 periods of less human impact (during the DA and the LIA)



# Some global implications...



A West – East Mediterranean dipole?



Thanks

**That's All!**

**Thanks for your attention**

**The study is funded by  
GLOBALKARST and GRACCIE  
CONSOLIDER-INGENIO projects  
(Spanish Government)**

**Thanks to the people that attended  
the coring expeditions and IPE  
technicians**

**Instituto Pirenaico  
de Ecología  
IPE-CSIC**

